

Quaternion Analyticity of Time-Harmonic Dyon Field Equations

Jivan Singh⁽¹⁾, P. S. Bisht⁽²⁾ and O. P. S. Negi⁽²⁾

February 2, 2008

⁽¹⁾Department of Physics
Govt. Post Graduate College
Pithoragarh- Uttarakhand, India

⁽²⁾Department of Physics
Kumaun University
S. S. J. Campus
Almora- 263601 (Uttarakhand), India

Email:-jgaria@indiatimes.com
ps_bisht123@rediffmail.com
ops_negi@yahoo.co.in

Abstract

Quaternion analysis of time dependent Maxwell's equations in presence of electric and magnetic charges has been developed in unique, simple and consistent manner. It has been shown that this theory is extended consistently to time-harmonic Maxwell's equation for dyons.

Dirac put forward the idea of magnetic monopole[1] while the fresh interests on the subject of monopoles was enhanced by the work of t' Hooft [2] and Polyakov [3] and its extension to the case of dyons by Julia and Zee [4] .

Consequently, these particles became intrinsic part of all current grand unified theories [5, 6] with their enormous potential importance [7, 8, 9, 10, 11, 12, 13] in connection with various physical problems. Keeping in view the results of Witten [13] that monopoles are necessarily dyons, a self-consistent and covariant quantum field theory for dyons each carrying the generalized charges as its real and imaginary parts has been constructed [14, 15] and accordingly the quaternionic forms of generalized field equations of dyons are developed [16, 2, 18, 19] in unique, simpler and compact notations. On the other hand Kravchenko [20] has analysed the Maxwell's equation in the presence of sources, time-dependent electromagnetic fields in homogeneous (isotropic) and chiral medium. In our previous papers, we [21, 22] have analysed the generalized Maxwell's -Dirac equations in homogenous(isotropic) medium, developed their quaternionic formulation and also obtained the solutions for the classical problem of moving dyons in simple, compact and consistent manner. Keeping in view all these facts in mind, in this paper, we have studied the Maxwell's equations in presence of electric and magnetic sources (i.e. dyons) and extended the generalised electromagnetic field equations associated with dyons to the case of time dependent harmonic Maxwell's-Dirac equations of dyons in simple, compact and consistent way. It has been emphasized that the theory reduces to the theory of dynamics of electric (magnetic) charge in the absence of magnetic (electric) charge on dyons.

Assuming the existence of magnetic monopoles in homogeneous (isotropic) medium, we have deduced the generalized Maxwell's-Dirac field equations of dyons in the following form [7];

$$\begin{aligned}
\vec{\nabla} \cdot \vec{E} &= \frac{\rho_e}{\epsilon} \\
\vec{\nabla} \cdot \vec{B} &= \mu \rho_m \\
\vec{\nabla} \times \vec{E} &= -\frac{\partial \vec{B}}{\partial t} - \frac{\vec{j}_m}{\epsilon} \\
\vec{\nabla} \times \vec{B} &= \frac{1}{v^2} \frac{\partial \vec{E}}{\partial t} + \mu \vec{j}_e
\end{aligned} \tag{1}$$

where ρ_e and ρ_m are respectively the electric and magnetic charge densities while $\vec{j}_e$ and $\vec{j}_m$ are the corresponding current densities, $\vec{E}$ is electric field intensity and $\vec{B}$ is magnetic field intensity while ϵ and μ are defined respectively as relative permittivity and permeability of the medium in electric and magnetic

fields. Differential equations (1) are the generalized field equations of dyons in homogeneous (isotropic) medium and the electric and magnetic fields are correspondingly called generalized electromagnetic fields of dyons. These electric and magnetic fields of dyons are expressed in terms of componets of two electromagnetic potentials in following differential form in homogeneous (isotropic) medium i.e.,

$$\vec{E} = -\vec{\nabla}\phi_e - \frac{\partial\vec{C}}{\partial t} - \vec{\nabla} \times \vec{D} \quad (2)$$

$$\vec{B} = -\vec{\nabla}\phi_m - \frac{1}{v^2}\frac{\partial\vec{D}}{\partial t} + \vec{\nabla} \times \vec{C} \quad (3)$$

where $\{C^\mu\} = \{\phi_e, v\vec{C}\}$ and $\{D^\mu\} = \{v\phi_m, \vec{D}\}$ are the four-potentials respectively associated with electric and magnetic charges of dyons. Let us define the complex vector field $\vec{\psi}$ in the following form [7]

$$\vec{\psi} = \vec{E} - iv\vec{B}. \quad (4)$$

Equations (2,3) and (4) then lead to the following relation between generalized field and the components of generalized four-potential as,

$$\vec{\psi} = -\frac{\partial\vec{V}}{\partial t} - \vec{\nabla}\phi - iv(\vec{\nabla} \times \vec{V}) \quad (5)$$

where $\{V_\mu\}$ is the generalized four-potential of dyons in homogeneous (isotropic) medium given by

$$V_\mu = \{\phi, \vec{V}\} \quad (6)$$

where

$$\phi = \phi_e - iv\phi_m \quad (7)$$

and

$$\vec{V} = \vec{C} - i \frac{\vec{D}}{v}. \quad (8)$$

Maxwell's field equation (1) in isotropic medium may then be written in terms of generalized field $\vec{\psi}$ as

$$\vec{\nabla} \cdot \vec{\psi} = \frac{\rho}{\epsilon} \quad (9)$$

$$\vec{\nabla} \times \vec{\psi} = -iv(\mu \vec{J} + \frac{1}{v^2} \frac{\partial \vec{\psi}}{\partial t}) \quad (10)$$

where ρ and $\vec{J}$ are the generalized charge and current source densities of dyons in homogeneous medium given by [7],

$$\rho = \rho_e - i \frac{\rho_m}{v} \quad (11)$$

$$\vec{J} = \vec{j}_e - iv \vec{j}_m. \quad (12)$$

In terms of complex potential the field equation is written as

$$\square \phi = v \mu \rho \quad (13)$$

$$\square \vec{V} = \mu \vec{J} \quad (14)$$

We may thus write the following tensorial form of generalized Maxwell's -Dirac equations of dyons in homogeneous (isotropic) medium [7] as

$$F_{\mu\nu,\nu} = j_\mu^e \quad (15)$$

$$F_{\mu\nu,\nu}^d = j_\mu^m. \quad (16)$$

Defining generalized field tensor of dyons as

$$G_{\mu\nu} = F_{\mu\nu} - iv F_{\mu\nu}^d. \quad (17)$$

One can directly obtain the following generalized field equation of dyons in homogeneous (isotropic) medium i.e.

$$G_{\mu\nu,\nu} = J_\mu. \quad (18)$$

Using the Fourier transform, any electromagnetic field can be represented as an infinite superposition of time-harmonic (monochromatic) fields. These fields are normally considered as the main object of study in radio electronics, wave propagation theory and many other branches of physics and engineering. A time harmonic electromagnetic field has the following form [20],

$$\vec{E}(x, t) = \text{Re}(\vec{E}(x)e^{-i\omega t}) \quad (19)$$

and

$$\vec{B}(x, t) = \text{Re}(\vec{B}(x)e^{-i\omega t}) \quad (20)$$

where the electric field $\vec{E}$ and magnetic field $\vec{B}$ depend on the spatial variables $x = (x_1, x_2, x_3)$ and all dependence on time is contained in the factor $e^{-i\omega t}$. $\vec{E}$ and $\vec{B}$ are expressed as the complex vectors called the complex amplitudes of electromagnetic field and ω denotes the frequency of oscillations. Substituting the values of $\vec{E}$ and $\vec{B}$ into the generalized dyonic equation (1) in isotropic medium, we obtain

$$\begin{aligned} \vec{\nabla} \cdot \vec{E} &= \frac{\rho_e}{\epsilon} \\ \vec{\nabla} \cdot \vec{B} &= \mu \rho_m \\ \vec{\nabla} \times \vec{E} &= -i\omega \vec{B} - \frac{\vec{j}_m}{\epsilon} \\ \vec{\nabla} \times \vec{B} &= -\frac{i\omega}{v^2} \vec{E} + \mu \vec{j}_e. \end{aligned} \quad (21)$$

Let us denote $\alpha = \omega \sqrt{\epsilon \mu} = \frac{\omega}{v}$, where the square root is chosen that $\text{Im} \alpha \geq 0$. The quantity α is called the wave number. Let us write the $\vec{D}$, $\vec{E}$ and $\vec{B}$ in the following quaternionic form as,

$$D = \partial_1 e_1 + \partial_2 e_2 + \partial_3 e_3 \quad (22)$$

$$E = E_1 e_1 + E_2 e_2 + E_3 e_3 \quad (23)$$

$$B = B_1 e_1 + B_2 e_2 + B_3 e_3 \quad (24)$$

where e_1, e_2 and e_3 are the elements of quaternion and satisfy the following multiplication rule,

$$\begin{aligned} e_0^2 &= 1 \\ e_j e_k &= -\delta_{jk} + \epsilon_{jkl} e_l \end{aligned} \quad (25)$$

where δ_{jk} and ϵ_{jkl} ($j, k, l = 1, 2, 3$ and $e_0 = 1$) are respectively the Kronecker delta and three-index Levi-Civita symbol. Using equations (22,23,24), we get the following quaternion differential equations i.e.

$$\begin{aligned} D \vec{E} &= (\partial_1 e_1 + \partial_2 e_2 + \partial_3 e_3)(E_1 e_1 + E_2 e_2 + E_3 e_3) \\ &= -\frac{\rho_e}{\epsilon} - \frac{\vec{j}_m}{\epsilon} + i\omega \vec{B} \end{aligned} \quad (26)$$

and

$$\begin{aligned} D \vec{B} &= (\partial_1 e_1 + \partial_2 e_2 + \partial_3 e_3)(B_1 e_1 + B_2 e_2 + B_3 e_3) \\ &= -\mu \rho_m + \mu \vec{j}_e - i \frac{\omega}{v^2} \vec{E}. \end{aligned} \quad (27)$$

Let us introduce the following pairs of purely vectorial biquaternionic functions,

$$\vec{l} = -\frac{i\omega}{v^2} \vec{E} + \alpha \vec{B} \quad (28)$$

and

$$\vec{m} = \frac{i\omega}{v^2}\vec{E} + \alpha\vec{B}. \quad (29)$$

Taking the divergence of third and fourth equation (21), we get the following pairs of continuity equation for electric and magnetic charges i.e.,

$$\vec{\nabla} \cdot \vec{j}_e - i\omega\rho_e = 0 \quad (30)$$

and

$$\vec{\nabla} \cdot \vec{j}_m - i\omega\mu\epsilon\rho_m = 0. \quad (31)$$

Applying the quaternionic operator D given by equation (22) to the quaternionic form of J the generalized current of dyons and using equations (26,27) and (30,31), we get

$$D\vec{l} = \frac{1}{\epsilon v^2}[\vec{\nabla} \cdot \vec{J}^*] + \alpha\mu\vec{J}^* + \alpha\vec{J} \quad (32)$$

where $\vec{J}^*$ is the complex conjugate of dyonic current density in homogeneous (isotropic) medium given by equation (12). Thus $\vec{J}$ satisfies the equation which is derived by equation (32) as,

$$(D - \alpha)\vec{J} = \mu[\vec{\nabla} \cdot \vec{J}^*] + \alpha\mu\vec{J}^*. \quad (33)$$

Analogous to equation (33), $\vec{m}$ also satisfies the equation (33) as,

$$(D + \alpha)\vec{m} = -\mu[\vec{\nabla} \cdot \vec{J}] + \alpha\mu\vec{J}. \quad (34)$$

Thus, the process of diagonalization can be written in the matrix form as

$$\begin{pmatrix} D & -i\omega \\ i\omega & D \end{pmatrix} \begin{pmatrix} \vec{E} \\ \vec{B} \end{pmatrix} = B_\alpha \begin{pmatrix} D - \alpha & 0 \\ 0 & D + \alpha \end{pmatrix} B_\alpha^{-1} \begin{pmatrix} \vec{E} \\ \vec{B} \end{pmatrix} \quad (35)$$

where

$$B_{\alpha} = \begin{pmatrix} -\frac{i\omega}{v^2} & \alpha \\ \frac{i\omega}{v^2} & \alpha \end{pmatrix} \quad (36)$$

and

$$B_{\alpha}^{-1} = \begin{pmatrix} -\frac{v^2}{i\omega} & \frac{v^2}{i\omega} \\ \frac{1}{\alpha} & \frac{1}{\alpha} \end{pmatrix}. \quad (37)$$

As such, we have obtained the two decoupled equations for the unknown vectors $\vec{J}$ and $\vec{m}$, which simplifies the analysis of the generalized Maxwell's-Dirac equation of dyons in homogeneous (isotropic) medium. These equations reduce to the theory of electric charge (magnetic) monopole predicted earlier by Kravchenko [20] in the absence of magnetic (electric charge) or vice versa.

References

- [1] Dirac, P. A .M, PRC. R. Soc. London Sec , **A 133** (1931),60.
- [2] t' Hooft, G, Nucl.Phys., **B79** (1974),276.
- [3] Polyakov, A. M., JEPT Lett, **20** (1974), 194.
- [4] Julia. B and Zee. A, Phys. Rev. , **D11** (1975), 2227.
- [5] Dokos C and Tomaros, T., Phy. Rev., **D21** (1980), 2940.
- [6] Preskill J. Annu. Rev. Nucl. Part. Sci., **34** (1984), 461.
- [7] Callen C. G., Phys. Rev., **D25** (1982), 2141.
- [8] Rubakov V.A., Nucl. Phy. B., **203** (1982), 211.
- [9] Mandelstam S., Phys. Rev., **D19** (1976), 249.
- [10] t' Hooft G., Nucl.Phys., **B138** (1978),1.
- [11] Rajput, B. S., J. Math, Phys. , **25** (1984), 351.
- [12] Rajput, B. S. and Rashmi Gunwant, Ind. J. of Pure and Appl. Phys., **26** (1988), 538.

- [13] Witten .E, Phy. Lett. , **B86** (1979), 283.
- [14] Rajput, B. S. and D. C. Joshi, Had. J., **4** (1981), 1805.
- [15] Rajput, B. S. and D. S. Bhakuni, Let. Nuovo Cimento, **34** (1982), 509.
- [16] Bisht, P.S., Negi, O.P.S and Rajput B.S., Int. J. Theor. Phys., **32** (1993), 2099.
- [2] Bisht, P.S., Negi, O.P.S and Rajput B.S., Il Nuovo Cimento, **A104** (1991), 337.
- [18] Bisht, P.S., Negi, O.P.S and Rajput B.S., Prog. Theor. Phys., **85** (1991), 151.
- [19] Bisht Shalini, Bisht, P.S. and Negi, O.P.S , Il Nuovo Cimento, **B113** (1998), 1449.
- [20] Kravchenko, V.V., Applied Quaternionic Analysis, Research and Exposition in Mathemetics, **28** (2003), Heldermann Verlage, Germany.
- [21] Jivan Singh, P. S. Bisht and O. P. S. Negi, “Generalized fields of dyons in Isotropic Medium”, arXiv:hep-th/0611208 .
- [22] Jivan Singh, P. S. Bisht and O. P. S. Negi, “Quaternion Analysis of generalized fields of dyons in Isotropic Medium”, arXiv:hep-th/0703083.